arXiv:1304.3356v1 [hep-ph] 11 Apr 2013

Chiral Symmetry Restoration and Scalar-Pseudoscalar partners in QCD

A. Gómez Nicola,[1, *] J. Ruiz de Elvira,[1, 2, †] and R. Torres Andrés[1, ‡]

[1]*Departamento de Física Teórica II. Univ. Complutense. 28040 Madrid. Spain.*
[2]*Helmholtz-Institut für Strahlen- und Kernphysik, Universität Bonn, D-53115 Bonn, Germany*

We present a consistent description of Scalar-Pseudoscalar partner degeneration at the QCD chiral transition in terms of the dominant low-energy physical states for the light quark sector. First, we obtain within model-independent one-loop Chiral Perturbation Theory (ChPT) that the QCD pseudoscalar susceptibility is proportional to the quark condensate. We also show that this chiral-restoring behaviour is compatible with recent lattice results for screening masses and gives rise to degeneration between the scalar and pseudoscalar susceptibilities (χ_S, χ_P) around the transition point, consistently with an $O(4)$-like current restoration pattern. This scenario is clearly confirmed by lattice data when we compare $\chi_S(T)$ with the quark condensate, expected to scale as $\chi_P(T)$. In addition, we show that saturating χ_S with the $\sigma/f_0(500)$ broad resonance observed in pion scattering and including its finite temperature dependence, account for the peak structure of $\chi_S(T)$ in lattice data and the associated critical temperature. This is carried out within a unitarized ChPT scheme which generates the resonant state dynamically and is also consistent with partner degeneration.

PACS numbers: 11.10.Wx, 11.30.Rd, 12.39.Fe, 12.38.Gc.

Chiral symmetry restoration $SU_V(N_f)\times SU_A(N_f)\to SU_V(N_f)$ with N_f light quark flavours has been a milestone in our present understanding of the Quantum Chromodynamics (QCD) phase diagram and Hadronic Physics under extreme conditions of temperature T and baryon density, as those produced in Heavy-Ion and Nuclear Matter experimental facilities such as RHIC, CERN (ALICE) and FAIR. Lattice simulations support that deconfinement and chiral restoration take place very close to one another in the phase diagram. In the physical case $N_f=2+1$ ($0\neq m_u=m_d\equiv m_q\ll m_s$) and for vanishing baryon chemical potential, they point towards a smooth crossover transition at pseudocritical temperature $T_c\sim$ 145-165 MeV [1, 2], the results being fairly consistent with the $O(4)$ universality class [3], which would hold for two light flavours in the chiral limit $m_q=0$. The crossover nature of the transition means in particular that there is no unique way to identify the transition point, the most efficient one in lattice being the scalar susceptibility peak position, rather than the vanishing point for the quark condensate $\langle\bar{q}q\rangle_T$, the order parameter, which decreases asymptotically with T for $m_q\neq 0$.

The equivalence with the $O(4)\to O(3)$ breaking pattern, which in its simplest linear realization takes place through the σ-component of the $O(4)$ field (σ,π^a) acquiring a thermal vacuum expectation value and mass vanishing at the transition in the chiral limit, led to early proposals of $\pi-\sigma$ meson degeneration ("chiral partners") at chiral restoration [4]. Degeneration in the vector-axial vector sector (ρ and a_1 states) as a signature of chiral restoration has also been thoroughly studied [5]. Nowadays, we know that the σ state is well established as a $\pi\pi$ scattering broad resonance for isospin and angular momentum $I=J=0$, known as $f_0(500)$ [6], which is then difficult to accommodate as an asymptotically free state. Therefore, in order to study chiral partner degeneration in the scalar-pseudoscalar sector, it is more appropriate to analyze the corresponding currents, which can be derived from a chiral effective lagrangian without introducing explicitly a particle-like σ degree of freedom. The scalar and pseudoscalar susceptibilities in terms of the corresponding QCD $SU(2)$ currents are given by:

$$\chi_S(T) \;=\; -\frac{\partial}{\partial m}\langle\bar{q}q\rangle_T=\int_E d^4x\left[\langle\mathcal{T}(\bar{q}q)(x)(\bar{q}q)(0)\rangle_T-\langle\bar{q}q\rangle_T^2\right]=\int_E d^4x\left[\frac{\delta}{\delta s(x)}\frac{\delta}{\delta s(0)}Z[s,p]\Big|_{s=m_q,p^a=0}\right], \quad (1)$$

$$\chi_P(T)\delta^{ab} \;=\; \int_E d^4x\langle\mathcal{T}P^a(x)P^b(0)\rangle_T=\int_E d^4x\left[\frac{\delta}{\delta p^a(x)}\frac{\delta}{\delta p^b(0)}Z[s,p]\Big|_{s=m_q,p^a=0}\right], \quad (2)$$

where $q=(u,d)$ is the quark field, the pseudoscalar current $P^a(x)=\bar{q}\gamma_5\tau^a q(x)$, the Euclidean four-volume $\int_E d^4x=\int_0^\beta d\tau\int d^3\vec{x}$ with $\beta=1/T$ and $\langle\cdot\rangle_T$ is a thermal average. For χ_P, Parity invariance of the QCD vacuum ($\langle P^a\rangle_T=0$) and isospin symmetry have been used. $Z[s,p]$ is the QCD generating functional with scalar and pseudoscalar sources coupled to the massless lagrangian in the light sector as $-s(x)(\bar{q}q)(x)+ip_a(x)P^a(x)$, so that $Z[m_q,0]$ is the QCD partition function.

Thus, should the scalar and pseudoscalar currents become degenerate at chiral restoration, $\chi_P(T)$ and $\chi_S(T)$ would meet at that point. The same problem has been studied in nuclear matter at $T=0$ [7] where it has been found that to leading order in nuclear density the pseudoscalar susceptibility decreases as the quark condensate. Therefore, since χ_S is expected to increase, as a measure of the fluctuations of the order parameter, at least up to the transition point, it seems plausible that they meet near the transition. In an ideal $O(4)$ pattern, the matching should take place at the maximum of χ_S.

Since P^a has the quantum numbers of the pion field π^a, its correlators are saturated by the pion state at low energies. Actually, from the low-energy theorems of current algebra (equivalent to the lowest order prediction of chiral lagrangians) one has $P^a \sim 2B_0F\pi^a$ (from PCAC theorem) with $B_0 = M^2/(2m_q)$ and F, M the pion decay constant and mass respectively, so that $\chi_P \sim 4B_0^2F^2G_\pi(p=0)$ with $G_\pi(p)$ the pion propagator. This is equivalent to saturate the correlator inside the integral in Eq.(2) with the dominant pion state. At this order one can take the free Euclidean pion propagator (interactions are suppressed at low energies) $G(p=0)=1/M^2$, so that using also the Gell-Mann-Oakes-Renner relation $M^2F^2 = -m_q\langle\bar{q}q\rangle$, we get $\chi_P \sim -\langle\bar{q}q\rangle/m_q$, as a first indication of the relation between the pseudoscalar susceptibility and the quark condensate. The latter result can actually be obtained formally as a Ward Identity (WI) from the QCD lagrangian [8], in connection with the definition of the quark condensate for lattice Wilson fermions [9]. However, both sides of the identity suffer from QCD renormalization ambiguities, so that this WI is formally well-defined only for exact chiral symmetry [9, 10]. It is therefore interesting to study how this identity is realized within Chiral Perturbation Theory (ChPT) [11], which describes the low-energy chiral symmetry broken phase of QCD in a model-independent framework. The previous current-algebra results are actually valid only at very low energies, i.e., to leading order (LO) in the ChPT expansion in powers of a generic scale p, denoting pion momenta or temperature, relative, respectively, to $\Lambda_\chi \sim 1$ GeV and T_c. In particular, the LO prediction for χ_P is temperature independent, so it is not obvious that it can be simply extrapolated as, say, $\langle\bar{q}q\rangle \to \langle\bar{q}q\rangle_T$. Actually, all the quantities involved are expected to change with temperature due to pion loop corrections, namely $M_\pi(T)$, $F_\pi(T)$ and $\langle\bar{q}q\rangle(T)$.

Similarly, from Eq.(1), one can relate χ_S with the propagator of a "σ-like state" such that it couples linearly to the external scalar source $s(x)$ in an explicit symmetry-breaking term $\mathcal{L}_{SB} = 2B_0Fs(x)\sigma(x)$. Without further specification about its nature and its coupling to other physical states such as pions, one already gets $\chi_S \sim 4B_0^2F^2G_\sigma(p=0)$, suggesting a growing behaviour inversely proportional to M_σ^2 as the sigma state reduces its mass to become degenerate with the pion.

The Next to Leading Order (NLO) corrections to χ_P can be obtained systematically and in a model-independent way within ChPT, where one can also calculate the scalar susceptibility χ_S to a given order only in terms of pion degrees of freedom. The price to pay is that we expect to reproduce only the behaviour of $\chi_{S,P}(T)$ for low and moderate temperatures. However, since χ_P is dominated by pions, whose dynamics are well described through ChPT, we expect to obtain a reasonable qualitative description of its T behaviour, whereas standard ChPT misses the peak structure of χ_S near the transition. We note in turn that the LO for χ_S vanishes, unlike that of χ_P. The ChPT NLO result for $\chi_S(T)$ can be found in [12, 13]. For $\chi_P(T)$ we consider the effective lagrangian $\mathcal{L}_2+\mathcal{L}_4+\ldots$, where $\mathcal{L}_{2n} = \mathcal{O}(p^{2n})$, including their dependence on the pseudoscalar source p^a as given in [11]. We follow similar steps as in [13, 14], now for the pseudoscalar correlator. The LO comes from $\mathcal{L}_2$ only and the NLO includes: a) G_π corrections to the LO, including one-loop $\mathcal{L}_2$ and tree level $\mathcal{L}_4$ terms, b) pion self-interactions $\mathcal{O}(\pi p^a \times \pi^3 p^b)$ in $\mathcal{L}_2$, c) crossed terms $\mathcal{L}_2 = \mathcal{O}(p^a\pi) \times \mathcal{L}_4 = \mathcal{O}(p^b\pi)$ and d) $\mathcal{L}_4 = \mathcal{O}(p^ap^b\pi^2)$ terms. The final result is:

$$\chi_P^{ChPT}(T) \;=\; 4B_0^2\left[\frac{F^2}{M^2}+\frac{1}{32\pi^2}(4\bar{h}_1-\bar{l}_3)-\frac{3}{2M^2}g_1(M,T)\right]+\mathcal{O}(F^{-2}) = -\frac{\langle\bar{q}q\rangle^{ChPT}(T)}{m_q}+\mathcal{O}(F^{-2}), \tag{3}$$

where subleading terms are labeled by their F^2 dependence, the first term inside brackets is the LO current-algebra $\mathcal{O}(F^2)$, F and M are the lagrangian pion mass and decay constant, related to the physical values $M_\pi \simeq 140$ MeV, $F_\pi \simeq 93$ MeV by $\mathcal{O}(F^{-2})$ corrections [11] and $\bar{l}_3$, $\bar{h}_1$ are renormalized scale-independent low-energy constants (LEC) of $\mathcal{L}_4$ [11] such that the result in Eq.(3) is finite and independent of the low-energy renormalization scale. The thermal loop function is given by:

$$g_1(M,T) = \frac{T^2}{2\pi^2}\int_{M/T}^{\infty} dx \frac{\sqrt{x^2-(M/T)^2}}{e^x-1}. \tag{4}$$



FIG. 1: Left: Comparison between the pseudoscalar screening mass ratio and $\Delta_{l,s}^{-1/2}$ for the lattice data in [16] (masses) and [17] (condensate) with the same lattice action and resolution and $T_c \simeq 196$ MeV. Right: Scalar susceptibility versus $\Delta_{l,s} \sim \chi_P(T)/\chi_P(0)$ from the data in [1] for which $T_c \simeq 155$ MeV. In both graphs $\Delta_{l,s} = r(T)/r(0)$ with $r = \langle\bar{q}q\rangle - (2m_q/m_s)\langle\bar{s}s\rangle$.

Therefore, we have obtained the WI connecting $\langle\bar{q}q\rangle$ and χ_P to NLO in model-independent ChPT, including finite-T effects. Furthermore, the m_q dependence cancels in $\chi_P(T)/\chi_P(0) = \langle\bar{q}q\rangle(T)/\langle\bar{q}q\rangle(0)$, where only meson parameters show up. To this order, $\chi_P(T)/\chi_P(0) = 1 - 3g_1(M,T)/(2F^2)$ so that the LEC dependence also disappears. Note that $\bar{h}_1$ comes from a contact term in $\mathcal{L}_4$ and is therefore another source of ambiguity in the NLO condensate [11].

We draw two important conclusions from the previous results. First, $\chi_P(T)$ has a critical nature, scaling like the order parameter instead of the much softer behaviour $\chi_P(T) \sim 1/M_\pi^2(T)$ with $M_\pi(T)$ the thermal pion mass, given at this order in [15], as a naive extension $M \to M_\pi(T)$ of the leading term would suggest. One can actually observe a clear signal of this critical behaviour in the lattice analysis of Euclidean correlators, which determine their large-distance space-like screening mass M^{sc} in different channels [16]. From Eq.(2) we expect $\chi_P = K_P(0) \sim (M_P^{pole})^{-2}$ with M_P^{pole} the pole mass associated to $K_P(p)$, the pseudoscalar correlator inside the integral in Eq.(2) in momentum space. Assuming a soft temperature behaviour for $M^{pole}(T)/M^{sc}(T)$, which is plausible below T_c, we can then explain the sudden increase of $M_P^{sc}(T)/M_P^{sc}(0)$ observed in this channel [16] since we expect that ratio to scale like $[\chi_P(0)/\chi_P(T)]^{1/2} \sim [\langle\bar{q}q\rangle(0)/\langle\bar{q}q\rangle(T)]^{1/2}$. As we show in Fig.1 (left panel) the correlation between these quantities is notorious, given the uncertainties involved. Data are taken from the same lattice group and under the same lattice conditions [16, 17]. Note that in lattice works $(2m_q/m_s)\langle\bar{s}s\rangle$ is subtracted from $\langle\bar{q}q\rangle$ in order to avoid renormalization ambiguities. Estimating the $T = 0$ condensates from NLO ChPT [21], this effect gives a 6% correction and, from the lattice values, it is about a 15% correction near T_c. Apart from the screening versus pole mass and the strange condensate corrections, one should not forget about the typical lattice uncertainties, like resolution, choice of action, staggered taste breaking and large pion masses [1, 17].



FIG. 2: Scalar versus pseudoscalar susceptibilities in ChPT and in our unitarized description. We show for comparison the lattice data of Fig.1 (right).

The second conclusion is that the decrease of χ_P and the increase of χ_S as they approach the critical point, leads to scalar-pseudoscalar susceptibility partner degeneration, which in an ideal $O(4)$ pattern should take place near the χ_S peak. Once again, this behaviour is observed in lattice data. In Fig.1 (right panel) we plot the subtracted condensate, expected to scale as $\chi_P(T)/\chi_P(0)$, versus $\chi_S(T)/\chi_P(0)$, both from the lattice analysis in [1]. The $T = 0$ values are taken from ChPT. The current degeneration is evident, not only at the critical point but also above it, where those two quantities remain very close to one another.

We now turn to our analytical results based on effective theories, which are plotted in Fig.2. ChPT to NLO gives an increasing $\chi_S(T)$, matching $\chi_P(T)$ at $T_d \simeq 0.9T_c$, where $\langle\bar{q}q\rangle^{ChPT}(T_c) = \chi_P^{ChPT}(T_c) = 0$. Although the ChPT expansion is not expected to reproduce the critical behaviour, this matching is a model-independent prediction. Near the chiral limit $M_\pi \ll T$, where critical effects are meant to be enhanced, the degeneration point $T_d = T_c - 3M_\pi/4\pi + \mathcal{O}(M_\pi^2/T_c)$, approaching the chiral restoration temperature in that limit.

Since the scalar susceptibility is dominated by the $I = J = 0$ lightest state, which does not show up in the ChPT expansion, let us consider its unitarized extension given by the Inverse Amplitude Method, which generates dynamically in $SU(2)$ the $f_0(500)$ and $\rho(770)$ resonances at $\vec{p} = \vec{0}$ as poles in the second Riemann sheet of the corresponding partial waves. More details and pole results at finite T can be found in [18, 19]. Saturating the scalar propagator with the $f_0(500)$ thermal state and assuming that its $p = 0$ mass does not vary much with respect to the pole mass, we have

$$\chi_S^U(T) = \frac{\chi_S^{ChPT}(0)M_S^2(0)}{M_S^2(T)}, \tag{5}$$

where we have normalized to the $T = 0$ ChPT value, since we are demanding that all our $T = 0$ results match the model-independent ChPT predictions. This normalization compensates partly the difference between the $p = 0$ and pole masses. Under this approximation, the full self-energy real part is $M_S^2 = M_p^2 - \Gamma_p^2/4$, where $(M_p - i\Gamma_p/2)^2$ is the pole position in p^2 [19] and the self-energy imaginary part vanishes at $p = 0$. Note that the width Γ_p for this state is not small compared to M_p.

We plot $\chi_S^U(T)$ in Fig.2. The result agrees with standard ChPT at low T and improves remarkably the behaviour near the transition. It develops a maximum at $T_c \simeq 157$ MeV, very much in agreement with the lattice data peak position. M_p is monotonically decreasing with T while Γ_p increases first due to thermal phase space but at some point mass reduction dominates and Γ_p decreases [19]. Furthermore, approaching the chiral limit by taking the $M_\pi = 10$ MeV poles from [19] gives a vanishing $M_S(T)$ at $T_c \simeq 118$ MeV and hence a divergent χ_S^U from Eq.(5) at T_c. Thus, we get the expected T_c reduction and stronger χ_S growth [3] near the chiral limit.

The quark condensate cannot be extracted directly from the unitarized susceptibility. However, we can obtain an approximate description by assuming that the relevant temperature and mass dependence, as far as the critical behaviour is concerned, comes from pion loop functions as $\delta\langle\bar{q}q\rangle^U(T,M) = B_0 T^2 g(T/M)$ and $\delta\chi_S = B_0^2 h(T/M)$, with $\delta f(T) = f(T) - f(0)$. This T/M dependence holds actually to NLO ChPT, as in Eq.(4). Then, from Eq. (1), since $\delta\chi_S = -\partial\delta\langle\bar{q}q\rangle/\partial m_q$, we get

$$g(x) = g(x_0) + \int_{x_0}^{x} \frac{h(y)}{y^3} dy \qquad \text{for } x > x_0, \tag{6}$$

with $T_0 = x_0 M \ll M$ a suitable low-T scale below which we use directly NLO ChPT, which has a better analytic behaviour near $T = 0$. The h function is obtained from the T dependence of χ_S in Eq.(5). The resulting $\langle\bar{q}q\rangle^U$ is plotted in Fig.2 with $T_0 \simeq 12$ MeV [22]. The critical behaviour is again nicely improved compared to the ChPT curves and is in better agreement with lattice data in that region. In addition, we obtain once more a scalar-pseudoscalar matching near the χ_S peak and hence of chiral restoration. The corresponding $\langle\bar{q}q\rangle^U$ near the chiral limit ($M_\pi = 10$ MeV) is much more abrupt, vanishing and matching χ_S at the same point, as expected.

Recall that in the above unitarized analysis, we are not performing a fit to lattice points. We just use the same LEC which generate the $T = 0$ physical $f_0(500)$, ρ states [19] and then provide our results for the susceptibility and condensate. Fitting the LEC within their uncertainties would improve the compatibility of our unitarized curves with lattice predictions below and near the transition. We do not expect our effective theory analysis to reproduce the chiral restoration pattern above T_c.

Summarizing, we have shown that a consistent scalar-pseudoscalar $O(4)$-like current degeneration pattern at chiral symmetry restoration emerges from lattice simulations and analytical effective theory analysis. The pseudoscalar susceptibility scales as the quark condensate and becomes degenerate with its "chiral partner" scalar susceptibility close to the scalar transition peak. The lattice data supporting this picture are well accounted for by the dominant physical states: pions and the $f_0(500)$ scalar resonance generated in pion scattering at finite temperature. Although we have restricted here, for simplicity, to the $N_f = 2$ case, the analysis can be extended to $N_f = 3$, where the role of other scalar states such as the $a_0(980)$ can also be studied [20].

Acknowledgments

Useful comments from F. Karsch, S. Mukherjee and D. Cabrera are acknowledged. Work partially supported by the EU FP7 HadronPhysics3 project, the Spanish project FPA2011-27853-C02-02 and FPI Programme (BES-2009-013672, R.T.A), and by the German DFG (SFB/TR 16, J.R.E.).

---

[*] Electronic address: gomez@fis.ucm.es
[†] Electronic address: elvira@hiskp.uni-bonn.de
[‡] Electronic address: rtandres@fis.ucm.es

[1] Y. Aoki *et al*, JHEP **0906**, 088 (2009).
[2] A. Bazavov *et al.*, Phys. Rev. D **85**, 054503 (2012).
[3] S. Ejiri *et al*, Phys. Rev. D **80**, 094505 (2009).
[4] T. Hatsuda and T. Kunihiro, Phys. Rev. Lett. **55**, 158 (1985).
[5] R. Rapp and J. Wambach, Adv. Nucl. Phys. **25**, 1 (2000).
[6] J. Beringer *et al.* [Particle Data Group Collaboration], Phys. Rev. D **86**, 010001 (2012) and references therein.
[7] G. Chanfray and M. Ericson, Eur. Phys. J. A **16**, 291 (2003).
[8] D. J. Broadhurst, Nucl. Phys. B **85**, 189 (1975).
[9] M. Bochicchio *et al*, Nucl. Phys. B **262**, 331 (1985).
[10] P. Boucaud *et al*, Phys. Rev. D **81**, 094504 (2010).
[11] J. Gasser and H. Leutwyler, Ann. Phys. **158**, 142 (1984).
[12] A. Gomez Nicola and R. Torres Andres, Phys. Rev. D **83**, 076005 (2011).
[13] A. Gomez Nicola, J. R. Pelaez and J. Ruiz de Elvira, Phys. Rev. D **87**, 016001 (2013).
[14] A. Gomez Nicola, J. R. Pelaez and J. Ruiz de Elvira, Phys. Rev. D **82**, 074012 (2010).
[15] J. Gasser and H. Leutwyler, Phys. Lett. B **184**, 83 (1987).
[16] M. Cheng *et al*, Eur. Phys. J. C **71**, 1564 (2011).
[17] A. Bazavov *et al*, Phys. Rev. D **80**, 014504 (2009).
[18] A. Dobado, A. Gomez Nicola, F. J. Llanes-Estrada and J. R. Pelaez, Phys. Rev. C **66**, 055201 (2002).
[19] D. Fernandez-Fraile, A. Gomez Nicola and E. T. Herruzo, Phys. Rev. D **76**, 085020 (2007).
[20] A. Gomez Nicola, J. Ruiz de Elvira and R. Torres Andres, work in preparation.

[21] For standard ChPT we use the same LEC values as in [12, 13]. Their influence is more important in χ_S, due to the vanishing of the LO, than in χ_P, $\langle\bar{q}q\rangle$.

[22] We find very small numerical differences changing T_0 between 10-60 MeV.